

Key to Genera of Lycidae of South Carolina

(adapted from Arnett 1960 and Green 1949-53)

- 1 Head rostrate; pronotum with a short longitudinal carina at apex of disc [*Lyconotus lateralis*](#) Melsheimer
Head not rostrate; pronotum various 2
- 2(1) Elytral costae feeble or indistinct, the fourth obliterated basally; intervals not reticulate
..... [*Calochromus perfacetus*](#) (Say)
Elytral costae well-defined; intervals reticulate 3
- 3(2) Pronotum with two or more longitudinal carinae on disc 4
Pronotum with or without a single longitudinal carina on disc 7
- 4(3) Longitudinal carinae of pronotum entire, confluent at base and apex, and enclosing a central diamond-shaped cell 5
Longitudinal carinae not entire, central cell not diamond-shaped 6
- 5(4) Elytron with four raised costae; reticulation in ten rows of cells [*Dictyoptera*](#)
Elytron with three raised carinae, reticulation in eight rows of cells [*Greenarus thoracicus*](#) (Randall)
- 6(4) Frons normally deflexed, not produced or transversely carinate; pronotum wider than long, sides convergent
from base to apex [*Eros humeralis*](#) (Fab.)
Frons acutely produced, transversely carinate between antennae; pronotum subrectangular, anterior half
of sides parallel [*Erotides sculptilis*](#) (Say)
- 7(3) Pronotum without carinae, with median longitudinal impression [*Plateros*](#) spp.
(identification of males based on genitalia)
Pronotum with at least a short carina on disc 8
- 8(7) Longitudinal carina of pronotum apical and short, oblique carina each side [*Eropterus*](#)
Longitudinal carina of pronotum entire 9
- 9(8) Antenna subserrate, compressed; elytra not broadly expanded [*Lopheros*](#)
Antenna markedly compressed, serrate or flabellate; elytra of many species broadly expanded 10
- 10(9) Apical maxillary palpomere large and markedly transverse; antennae serrate in both sexes [*Calopteron*](#)
Apical maxillary palpomere longer than wide; antennae flabellate in male, serrate in female 11
- 11(10) Apical labial palpomere transverse, broadly triangular, apically minutely digitate; mesosternum with entire
longitudinal median impression; elytra broadly expanded, each with four discal costae; intervals with
double row of cells [*Caenia dimidiata*](#) (Fab.)
Apical labial palpomere longer than wide; mesosternum not impressed; elytra slender, each with three discal
costae; intervals with single row of cells [*Leptoceletes basalis*](#) (LeConte)

Arnett, R. H., Jr. 1060. The beetles of the United States (a manual for identification). Catholic University of America Press, Washington, D. C. 1112 pp.

Green, J. W. 1949. The Lycidae of the United States and Canada, I. The tribe Lycini (Coleoptera). Transactions of the American Entomological Society. 75:53-70.

Green, J. W. 1950. The Lycidae of the United States and Canada, II. The tribe Lygostopterini (Coleoptera). Transactions of the American Entomological Society 76:13-25.

Green, J. W. 1951. The Lycidae of the United States and Canada, III. The tribe Platerodini (in part) (Coleoptera). Transactions of the American Entomological Society 77:1-20.

Green, J. W. 1952. The Lycidae of the United States and Canada, IV. The tribe Calopterini (Coleoptera). Transactions of the American Entomological Society 78:1-19.

Green, J. W. 1953. The Lycidae of the United States and Canada, V. *Plateros* (Coleoptera). Transactions of the American Entomological Society 78:149-181.

Kazantsev, S. V. 2004. Phylogeny of the tribe Erotini (Coleoptera, Lycidae), with descriptions of new taxa. Zootaxa 496:1-48.